Borghi, B. & Casati, A. Incidence and risk factors for allogenic blood transfusion during major joint replacement using an integrated autotransfusion regimen, 411
Borghi, B. *see* Casati, A.
Borghi, B. *see* Magistris, L.
Boris-Möller, F. *see* Fletcher, R.
Borissov, B. *see* Park, G.R.
Bouaggad, A., Al Harrar, R., Bouderka, M.A. & Abassi, O. The effect of oral etilefrine premedication on the incidence of hypotension during spinal anaesthesia, 177
Bouderka, M.A. *see* Bouaggad, A.
Bovill, J.G. Opioid detoxification under anaesthesia, 657
Bovill, J.G. *see* El-Kerdawy, H.M.M.
Bovill, J.G. *see* Novak-Jankovič, V.
Bozkurt, P. *see* Altintaş, F.
Brill, G. *see* Brill, S.
Brill, S., Gurman, G.M. & Brill, G. Evaluation of the VersaMed portable ventilator: clinical trials, 737
Brimacombe, J. *see* Agrò, F.
Brimacombe, J. *see* Kihara, S.
Brodner, G., Mertes, N., Buerkle, H., Marcus, M.A.E. & Van Aken, H. Acute pain management: analysis, implications and consequences after prospective experience with 6349 surgical patients, 566
Brodner, G., Scherer, R. & Van Aken, H. Continuous spinal anaesthesia or continuous epidural anaesthesia for postoperative pain control after hip replacement, 402
Brouwers, M.J.A., Driessen, J.J. & Severijnen, R.S.V.M. Epidural analgesia in a newborn with Hirschsprungís disease, associated with congenital central hypoventilation syndrome, 751
Brown, J. *see* Moss, H.A.
Brown, R.A. *see* Philip, B.K.
Browne, J., Fung, M., Donnelly, M. & Cooney, C. The use of EMLA reduces the pain associated with digital ring block for ingrowing toenail correction, 182
Buerkle, H. *see* Brodner, G.
Buerkle, H., Huge, V., Wolfgart, M., Steinbeck, J., Mertes, N., Van Aken, H. & Prien, T. Intra-articular clonidine analgesia after knee arthroscopy, 295
Bugge, J.F. & Tanbo, T. Nitric oxide in the treatment of fulminant pulmonary failure in a young pregnant woman with varicella pneumonia, 269
Buhre, W. *see* Kazmaier, S.
Bui-Xuan, B. *see* Tabib, A.
Busch, T. *see* Kazmaier, S.
Buscher, H. *see* Kazmaier, S.

Çakan, T. *see* Ceyhan, A.
Calam, R.M., Jimmieson, P., Cox, A.D., Glasgow, D.V., Groth, S. & Larsen, S. Can computer-based assessment help us understand children's pain?, 284
Campbell, P. *see* Jaffe, S.M.
Caria, M.A. *see* De Riu, P.L.
Casati, A. *see* Borghi, B.
Casati, A. *see* Magistris, L.
Casati, A., Comotti, L., Tommasino, C., Leggieri, C., Bignami, E., Tarantino, F. & Torri, G. Effects of pneumoperitoneum and reverse Trendelenburg position on cardiopulmonary function in morbidly obese patients receiving laparoscopic gastric banding, 300
Casati, A., Gallioli, G., Scandroglio, M., Passaretta, R., Borghi, B. & Torri, G. Accuracy of end-tidal carbon dioxide monitoring using the NBP-75® microstream capnometer. A study in intubated ventilated and spontaneously breathing non-intubated patients, 622
Celich, M.F. *see* Peixoto, A.J.
Cevheroglu, D. *see* Bilgin, H.
Ceyhan, A., Gülhan, Y., Çakan, T., Adişen, Y., Turgut, Ö. & Ünal, N. Anaesthesia for Proteus syndrome, 645
Chakrabarti, M.K. *see* Pac-Soo, C.K.
Chakrabarti, M.K. *see* Whitwam, J.G.
Chan, M.T.V. & Gin, T. What does the bispectral EEG index monitor?, 146
Chanimov, M., Berman, S., Weissgarten, J., Averbukh, Z., Cohen, M., Grinshpun, Y. & Bahar, M. Substances used for local and general anaesthesia in major surgery suppress proliferative responsiveness of normal rat peripheral blood mononuclear cells in culture, 248
Chatzimichali, A. *see* Michaloudis, D.

Chelly, J.E. *see* Philip, B.K.
Chevalley, C., Montessuit, M., Murith, N. & Licker, M. Near-fatal acute dissection of the thoracic aorta involving the right coronary artery, 709
Chippa, J.H. *see* Popat, M.T.
Cohen, C. *see* Didailler, J.L.
Cohen, M. *see* Chanimov, M.
Coley, S. *see* Shah, J.
Comotti, L. *see* Casati, A.
Connolly, J. & Cunningham, A.J. Preoperative fasting and administration of regular medications in adult patients presenting for elective surgery. Has new evidence changed clinical practice?, 219
Cooke, T. *see* Lydon, A.
Cooney, C. *see* Browne, J.
Cox, A.D. *see* Calam, R.M.
Cross, M.H. *see* O'Connor, P.J.
Crozier, T. The anaesthetic potency of propofol in the rat, 276
Cruaud, P. *see* Asehnoune, K.
Cunningham, A.J. *see* Connolly, J.
Cvachovec, K., Horáček, M. & Vislocký, I. A retrospective survey of fibrinolysis as an indicator of poor outcome after cardiopulmonary bypass and a possible early sign of systemic inflammation syndrome, 173

Dahl, O.E. Thromboembolism and thromboprophylaxis in high risk surgery: facts and assumptions – a topic for emotions?, 343
Danelli, G. *see* Magistris, L.
Davies, M.W. *see* Moore, E.W.
De La Cuesta, F. *see* Avellaneda, C.
De Lange, J.J. *see* El Azab, S.R.
De Mey, J.C., Strobbe, J., Poelaert, J., Hoebeke, P. & Mortier, E. The influence of sufentanil and/or clonidine on the duration of analgesia after a caudal block for hypospadias repair surgery in children, 379
De Riu, G. *see* De Riu, P.L.
De Riu, P.L., De Riu, G., Testa, C., Mulas, M., Caria, M.A., Mameli, S. & Mameli, O. Disposition of propofol between red blood cells, plasma, brain and cerebrospinal fluid in rabbits, 18
De Rossi, L., Fritz, H. & Klein, U. Comparison of cisatracurium-induced neuromuscular block at the masseter and adductor pollicis muscle, 583
De Wolff, M.H., Leather, H. & Wouters, A. Effect of clevidipine, an ultra-short acting 1,4-dihydropyridine calcium channel blocking drug, on the potency of isoflurane in rats and dogs, 506
Demian, A.D., Esmail, O.M. & Atallah, M.M. Acid-base equilibrium during capnoretroperitoneoscopic nephrectomy in patients with end-stage renal failure: a preliminary report, 256
Deni, F. *see* Magistris, L.
Dershwitz, M. *see* Philip, B.K.
Dib, M. *see* Didailler, J.L.
Didailler, J.L., Dominici, L., Dib, M., Cohen, C. & The Study Group Of Ketoprofen. The analgesic efficacy and tolerance of ketoprofen (100 mg) combined with morphine in patient-controlled analgesia after orthopaedic surgery, 459
Dimitriou, V. & Voyagis, G.S. Use of an illuminated flexible catheter for light-guided tracheal intubation through the intubating laryngeal mask by nurses, 46
Dimitriou, V. & Voyagis, G.S. Use of the intubating laryngeal mask for airway management and light-guided tracheal intubation in the lateral position, 395
Dimitriou, V. *see* Voyagis, G.S.
Dinter, W. *see* Schlack, W.
Dirksen, R. *see* Slappendel, R.
Dominici, L. *see* Didailler, J.L.
Donnelly, M. *see* Browne, J.
Dorn, C. *see* Krenn, H.
Dort, J.P. *see* Pendeville, P.E.
Driessen, J.J. *see* Brouwers, M.J.A.
Duggan, P.F. *see* Halpenny, M.
Duggan, P.F. *see* Lydon, A.
Dumont, L. & Mardirosoff, C. Effects of a pneuperitoneum in the obese patient, 786
Dunkel, M. *see* St Pierre, M.

El Azab, S.R., Scheffer, G.J., Rosseel, P.M.J. & De Lange, J.J. Induction and maintenance of anaesthesia with sevoflurane in comparison to high dose opioid during coronary artery bypass surgery, 336

El-Kerdawy, H.M.M., Van Zalingen, E.E. & Bovill, J.G. The influence of the α_2- adrenoceptor agonist, clonidine, on the EEG and on the MAC of isoflurane, 105
Elliott, T.S.J. *see* Moss, H.A.
Enel, D. *see* Gentili, M.E.
Esmail, O.M. *see* Demian, A.D.
Evans, T.N. *see* Park, G.R.

Fanelli, G. *see* Magistris, L.
Fanning, N.F. *see* Goto, Y.
Faroqui, M.H. *see* Moss, H.A.
Fernandez, C. *see* Gomar, C.
Filipovic, M. *see* Steiner, L.A.
Fitzgerald, R.D. *see* Krenn, H.
Fletcher, I.R. Clinical governance, 471
Fletcher, R. & Boris-Möller, F. Can we improve the estimate of arterial PCO_2 from end-tidal PCO_2?, 306
Flisberg, P., Törnebrandt, K., Walther, B. & Lundberg, J. A comparison of the effects on post-operative pain relief of epidural analgesia started before or after surgery, 627
Flossos, A. *see* Michaloudis, D.
Flossos, A. *see* Petru, A.
Flucker, C.J.R., Hart, E., Weisz, M., Griffiths, R. & Ruth, M. The 50-millilitre syringe as an inexpensive training aid in the application of cricoid pressure, 443
Frerichs, A. *see* Klockgether-Radke, A.P.
Fritz, H. *see* De Rossi, L.
Fujii, Y., Saitoh, Y., Tanaka, H. & Toyooka, H. Granisetron/dexamethasone combination for the prevention of postoperative nausea and vomiting after laparoscopic cholecystectomy, 64
Fung, M. *see* Browne, J.
Fung, M.A. *see* Gopinathan, C.

Gaffney, E. *see* Halpenny, M.
Gallagher, G. *see* Rae, C.P.
Galle-Röd, A. *see* Azad, S.C.
Galletly, D.C. *see* Whitwam, J.G.
Gallioli, G. *see* Casati, A.
Garcia-Guasch, R. *see* Lucas, M.
Gentili, M.E., Enel, D. & Blamoutier, A. Successful epidural blood patch in a patient with headache for 6 months after lumbar root decompression, 208
Giesecke, K. *see* Reihnér, E.
Gin, T. *see* Chan, M.T.V.
Giorgi, E. *see* Agostoni, M.
Giorgini, C. *see* Peduto, V.A.
Glasgow, D.V. *see* Calam, R.M.
Gockner, G. *see* Ihra, G.
Gomar, C. & Fernandez, C. Epidural analgesia-anaesthesia in obstetrics, 542
Gómez, A. *see* Avellaneda, C.
Gopinathan, C., Sockalingham, I., Fung, M.A., Peat, S. & Hanna, M.H. A comparative study of patient-controlled epidural diamorphine, subcutaneous diamorphine and an epidural diamorphine/bupivacaine combination for postoperative pain, 189
Gorce, P. *see* Asehnoune, K.
Goren, S. *see* Kahveci, S.F.
Goto, Y., Ho, S.L., McAdoo, J., Fanning, N.F., Wang, J., Redmond, H.P. & Shorten, G.D. General versus regional anaesthesia for cataract surgery: effects on neutrophil apoptosis and the postoperative pro-inflammatory state, 474
Griffiths, R. *see* Flucker, C.J.R.
Grinshpun, Y. *see* Chanimov, M.
Groeben, H., Großwendt, T., Silvanus, M.-T., Beste, M. & Peters, J. Lidocaine inhalation for local anaesthesia and attenuation of bronchial hyperreactivity with least airway irritation. Effect of three different dose regimens, 672
Groop, P.-H. *see* Norio, K.
Großwendt, T. *see* Groeben, H.
Groth, S. *see* Calam, R.M.
Grunditz, R. *see* Reihnér, E.
Guard, B.C. *see* Milner, Q.J.W.
Gülhan, Y. *see* Ceyhan, A.
Gunning, K.E. *see* Park, G.R.
Gurman, G.M. *see* Brill, S.
Gustafsson, L.L. *see* Reihnér, E.

Haemmerle, A. *see* Klockgether-Radke, A.P.
Hahn, R.G. & Löfgren, A. Epinephrine, potassium

and the electrocardiogram during regional anaesthesia, 132
Hahn, R.G. The volumetric fluid balance as a measure of fluid absorption during transurethral resection of the prostate, 559
Hahne, W.F. *see* Philip, B.K.
Halpenny, M., Markos, F., Snow, H.M., Duggan, P.F., Gaffney, E., O'Connell, D.P. & Shorten, G.D. The effects of fenoldopam on renal blood flow and tubular function during aortic cross-clamping in anaesthetized dogs, 491
Hammond, E.J. & Sweeney, B.P. Electronic data collection by trainee anaesthetists using palm top computers, 91
Hanna, M.H. *see* Gopinathan, C.
Hart, B. 'Diprivan': a change of formulation, 71
Hart, E. *see* Flucker, C.J.R.
Hay, H. Contamination of piped medical gas supply with water, 512
Hay, H. Delivery of an hypoxic gas mixture due to a defective rubber seal of a flowmeter control tube, 456
Hay, H. Oxygen delivery failure resulting from interference with a Bain breathing system, 591
Hay, H. Rapidly developing airway obstruction resulting from achalasia of the oesophagus, 398
Hein, L. *see* Nielsen, M.B.
Hellige, G. *see* Klockgether-Radke, A.P.
Hellige, G. *see* Klockgether-Radke, A.P.
Henderson, J.J. Questions about the Macintosh laryngoscope and technique of laryngoscopy, 2
Henderson, K. *see* Skelton, V.A.
Henderson, K.A. & Matthews, I.P. Staff exposure to anaesthetic gases in theatre and non theatre areas, 149
Hepağuşlar, H. *see* Aysel, İ.
Herbst, T. *see* Moss, H.A.
Hering, W. *see* St Pierre, M.
Hidestål, J. *see* Bodelsson, G.
Higashi, M. *see* Yamaura, K.
Hinojosa, M. *see* Lucas, M.
Hirschi, M., Meistelman, C. & Longrois, D. Effects of normothermic cardiopulmonary bypass on bispectral index, 499
Ho, S.L. *see* Goto, Y.
Hoebeke, P. *see* De Mey, J.C.
Hoermann, C. *see* Kuenszberg, E.
Hooper, R.H. *see* Jones, P.R.M.
Hopkins, A. *see* O'Riordan, J.
Horáček, M. *see* Cvachovec, K.
Hübler, M., Litz, R.J. & Albrecht, D.M. Combination of balanced and regional anaesthesia for minimally invasive surgery in a patient with myasthenia gravis, 325
Huge, V. *see* Buerkle, H.
Hutchins, J. *see* Park, G.R.
Hüttemann, E. *see* Meisner, M.

Ihan, A. *see* Novak-Jankovič, V.
Ihra, G., Gockner, G., Kashanipour, A. & Aloy, A. High-frequency jet ventilation in European and North American institutions: developments and clinical practice, 418
Inoue, H. *see* Yamaura, K.
Isaac, J.L. *see* Moss, H.A.
Isoniemi, H. *see* Norio, K.
Itonaga, Y. *see* Yamaura, K.

Jackson, A. *see* Watson, N.A.
Jaffe, S.M., Campbell, P., Bellman, M. & Baildam, A. Postoperative nausea and vomiting in women following breast surgery: an audit, 261
Jannopoulos, A. *see* Petru, A.
Jaszczyk, G. *see* Pakulski, C.
Jespersen, T.W. *see* Nielsen, M.B.
Jimmieson, P. *see* Calam, R.M.
Jones, P.R.M., Hooper, R.H., Stopp, N. & Moore, N.A. Nitric oxide delivery to the lung: a model, 160

Kızılarslan, S., Kuvaki, B., Onat, Ü. & Sağıroğlu, E. Epidural fentanyl–bupivacaine compared with clonidine–bupivacaine for analgesia in labour, 692
Kafkalaki, K. *see* Michaloudis, D.
Kahveci, S.F., Goren, S., Kutlay, O., Ozcan, B. & Korfali, G. Bedside percutaneous tracheostomy experience with 72 critically ill patients, 688
Kasakov, L. *see* Meisner, M.

Kashanipour, A. *see* Ihra, G.
Kassner, A. *see* Watson, N.A.
Katoh, T. *see* Suzuki, A.
Kaya, G. *see* Altintaş, F.
Kazmaier, S., Rathgeber, J., Buhre, W., Buscher, H., Busch, T., Mensching, K. & Sonntag, H. Comparison of ventilatory and haemodynamic effects of BIPAP and S-IMV/PSV for postoperative short-term ventilation in patients after coronary artery bypass grafting, 601
Keller, C. *see* Agrò, F.
Kettler, D. & Wilkinson, D. Confederation of European National Societies of Anaesthesiology (CENSA), 145
Kettler, D. *see* Klockgether-Radke, A.P.
Kettler, D. *see* Klockgether-Radke, A.P.
Khan, F. & Khan, F. A rare misplacement of a central venous catheter, 527
Khan, F. *see* Khan, F.
Kihara, S., Yaguchi, Y., Watanabe, S., Brimacombe, J., Taguchi, N. & Yamasaki, Y. Haemodynamic responses to the intubating laryngeal mask and timing of removal, 744
Kinsella, J. *see* Rae, C.P.
Kitajima, T. *see* Okuda, Y.
Klein, U. *see* De Rossi, L.
Kleinsasser, A. *see* Kuenszberg, E.
Klemola, U.-M. The psychology of human error revisited, 401
Klinck, J.R. *see* Park, G.R.
Klockgether-Radke, A.P., Frerichs, A., Kettler, D. & Hellige, G. Propofol and thiopental attenuate the contractile response to vasoconstrictors in human and porcine coronary artery segments, 485
Klockgether-Radke, A.P., Haemmerle, A., Kettler, D. & Hellige, G. Do muscle relaxants influence vascular tone in isolated coronary artery segments?, 481
Korfali, G. *see* Bilgin, H.
Korfali, G. *see* Kahveci, S.F.
Köszegi, T. *see* Molnár, Z.
Kovac, A. L. *see* Philip, B.K.
Krause, T. *see* Martens, U.
Krenn, H., Dorn, C., Fitzgerald, R.D., Pojer, H., Rumpold-Seitlinger, G. & List, W.F. Prognostic value of Apache II and III for preoperative evaluation of emergency surgery patients, 529
Kuenszberg, E., Loeckinger, A., Kleinsasser, A., Lindner, K.H., Puehringer, F. & Hoermann, C. Sevoflurane progressively prolongs the QT interval in unpremedicated female adults, 662
Kutlay, O. *see* Bilgin, H.
Kutlay, O. *see* Kahveci, S.F.
Kuvaki, B. *see* Kızılarslan, S.

Larsen, J.R. *see* Nielsen, M.B.
Larsen, S. *see* Calam, R.M.
Lassner, J. Who should provide anaesthesia?, 648
Lawton, G. *see* Weigert, A.
Leães, L.F. *see* Peixoto, A.J.
Leary, T.S., Milner, Q.J.W. & Niblett, D.J. The accuracy of the estimation of body weight and height in the intensive care unit, 698
Leather, H. *see* De Wolff, M.H.
Leggieri, C. *see* Casati, A.
Leonard, S.A., Walsh, M., Lydon, A., O'Hare, B. & Shorten, G.D. Evaluation of the effects of levobupivacaine on clotting and fibrinolysis using thromboelastography, 373
Licker, M. *see* Chevalley, C.
Lindgren, L. *see* Norio, K.
Lindner, K.H. *see* Kuenszberg, E.
List, W.F. *see* Krenn, H.
Litz, R.J. *see* Hübler, M.
Liu, C. *see* Skelton, V.A.
Loeckinger, A. *see* Kuenszberg, E.
Löfgren, A. *see* Hahn, R.G.
Lohs, T. *see* Meisner, M.
Loire, R. *see* Tabib, A.
Longrois, D. *see* Hirschi, M.
Lucas, M., Hinojosa, M., Rodríguez, A. & Garcia-Guasch, R. Anaesthesia in lipid myopathy, 461
Luis, M. *see* Santos, P.
Lund, N. *see* Vallet, B.
Lundberg, J. *see* Flisberg, P.
Lydon, A. *see* Leonard, S.A.
Lydon, A., Murray, C., Cooke, T., Duggan, P.F., O'Halloran, D. & Shorten, G.D. Evaluation of standard haemodynamic tests of autonomic function and HbA1c as predictors of delayed

gastric emptying in patients with type 1 diabetes mellitus, 99

Ma, D. *see* Whitwam, J.G.
Magistris, L., Casati, A., Albertin, A., Deni, F., Danelli, G., Borghi, B. & Fanelli, G. Combined sciatic–femoral nerve block with 0.75% ropivacaine: effects of adding a systemically inactive dose of fentanyl, 348
Mäkisalo, H. *see* Norio, K.
Malhotra, S.K. *see* Podder, S.
Malicier, D. *see* Tabib, A.
Mameli, O. *see* De Riu, P.L.
Mameli, S. *see* De Riu, P.L.
Marcus, M.A.E. *see* Brodner, G.
Mardirosoff, C. *see* Dumont, L.
Markos, F. *see* Halpenny, M.
Martens, U., Krause, T., Scholz, J., Wappler, F., Steinrücke, K. & Schulte Am Esch, J. Inositol 1,4,5-trisphosphate synthesis in mononuclear white blood cells of malignant hyperthermia-susceptible and normal human beings, following in vitro exposure to halothane, caffeine and ryanodine, 364
Martínez, G. *see* Santos, P.
Martos, F. *see* Avellaneda, C.
Matthews, I.P. *see* Henderson, K.A.
McAdoo, J. *see* Goto, Y.
McAnulty, G. *see* Snelgrove, H.
McKenzie, R. *see* Philip, B.K.
Meier-Hellmann, A. *see* Reinhart, K.
Meisner, M., Hüttemann, E., Lohs, T., Kasakov, L. & Reinhart, K. Elimination of procalcitonin and plasma concentrations during continuous veno-venous haemodiafiltration in septic patients, 665
Meistelman, C. *see* Hirschi, M.
Mensching, K. *see* Kazmaier, S.
Mertes, N. *see* Brodner, G.
Mertes, N. *see* Buerkle, H.
Mezzetti, D. *see* Peduto, V.A.
Michaloudis, D. *see* Petru, A.
Michaloudis, D., Petrou, A., Bakos, P., Chatzimichali, A., Kafkalaki, K., Papaioannou, A., Zeaki, M. & Flossos, A. Continuous spinal anaesthesia/analgesia for the perioperative management of high-risk patients, 239
Miller, D.M. Modification of the Penlon Nuffield 200 series ventilator and IPPV with a laryngeal mask airway, 354
Milliet, N. Recurarization in the recovery room, 332
Milner, Q.J.W. *see* Leary, T.S.
Milner, Q.J.W., Guard, B.C. & Allen, J.G. Alkalinization of amide local anaesthetics by addition of 1% sodium bicarbonate solution, 38
Mingus, M. *see* Philip, B.K.
Miras, A. *see* Tabib, A.
Møller, A. *see* Smith, A.
Molnár, Z., Szakmány, T., Köszegi, T. & Tekeres, M. Microalbuminuria and serum procalcitonin levels following oesophagectomy, 464
Monk, T.G. *see* Philip, B.K.
Montessuit, M. *see* Chevalley, C.
Moore, E.W. & Davies, M.W. Inhalational versus intravenous induction. A survey of emergency anaesthetic practice in the United Kingdom, 33
Moore, N.A. *see* Jones, P.R.M.
Mortier, E. *see* De Mey, J.C.
Moss, H.A., Tebbs, S.E., Faroqui, M.H., Herbst, T., Isaac, J.L., Brown, J. & Elliott, T.S.J. A central venous catheter coated with benzalkonium chloride for the prevention of catheter-related microbial colonization, 680
Mulas, M. *see* De Riu, P.L.
Murith, N. *see* Chevalley, C.
Murray, C. *see* Lydon, A.

Niblett, D.J. *see* Leary, T.S.
Nielsen, M.B., Jespersen, T.W., Larsen, J.R. & Hein, L. The effects of atropine on dynamic compliance in healthy intubated adults, 236
Norio, K., Mäkisalo, H., Isoniemi, H., Groop, P.-H., Pere, P. & Lindgren, L. Are diabetic patients in danger at renal transplantation? An invasive perioperative study, 729
Novak-Jankovič, V., Paver-Eržen, V., Bovill, J.G., Ihan, A. & Osredkar, J. Effect of epidural and intravenous clonidine on the neuro-endocrine and immune stress response in patients undergoing lung surgery, 50

O'Connell, D.P. *see* Halpenny, M.
O'Connor, P.J., Welsh, K.R., Cross, M.H. & Shah, M.V. Cardiovascular responses to pulmonary artery catheterization, 168
O'Halloran, D. *see* Lydon, A.
O'Hare, B. *see* Leonard, S.A.
O'Riordan, J., Hopkins, A., Ravenscroft, P.M. & Stevens, A. Patient-controlled analgesia and urinary retention following lower limb joint replacement: prospective audit and logistic regression analysis, 431
Okuda, Y., Sato, H., Kitajima, T. & Asai, T. Airway obstruction during general anaesthesia in a child with congenital tracheomalacia, 642
Onat, Ü. *see* Kızılarslan, S.
Osredkar, J. *see* Novak-Jankovič, V.
Owen-Smith, O. *see* Shah, J.
Ozcan, B. *see* Kahveci, S.F.

Pac-Soo, C.K., Wang, C., Chakrabarti, M.K. & Whitwam, J.G. Comparison of the effects of inhalational anaesthetic agents on sympathetic activity in rabbits, 311
Pakulski, C., Świniarski, A. & Jaszczyk, G. High subarachnoid block for severe bronchospasm, 594
Papaioannou, A. *see* Michaloudis, D.
Paries, J. *see* Asehnoune, K.
Park, G.R., Evans, T.N., Hutchins, J., Borissov, B., Gunning, K.E. & Klinck, J.R. Reducing the demand for admission to intensive care after major abdominal surgery by a change in anaesthetic practice and the use of remifentanil, 111
Passaretta, R. *see* Casati, A.
Pattnaik, S.K. & Bodra, R. Ballotability of cuff to confirm the correct intratracheal position of the endotracheal tube in the intensive care unit, 587
Paver-Eržen, V. *see* Novak-Jankovič, V.
Payne, J.P. Bradycardia, hypotension and cerebral dysfunction after spinal anaesthesia, 334
Pearman, M.H. *see* Philip, B.K.
Peat, S. *see* Gopinathan, C.
Pedersen, T. *see* Smith, A.
Peduto, V.A., Mezzetti, D., Properzi, M. & Giorgini, C. Sevoflurane provides better recovery than propofol plus fentanyl in anaesthesia for day-care surgery, 138
Peixoto Filho, A.J. *see* Peixoto, A.J.
Peixoto, A.J., Peixoto Filho, A.J., Leães, L.F., Celich, M.F. & Barros, M.A.V. Efficacy of prophylactic droperidol, ondansetron or both in the prevention of postoperative nausea and vomiting in major gynaecological surgery. A prospective, randomized, double-blind clinical trial, 611
Pendeville, P.E., Von Montigny, S., Dort, J.P. & Veyckemans, F. Double-blind randomized study of tramadol vs. paracetamol in analgesia after day-case tonsillectomy in children, 576
Pere, P. *see* Norio, K.
Peter, K. *see* Azad, S.C.
Peters, J. *see* Groeben, H.
Petrou, A. *see* Michaloudis, D.
Petru, A., Michaloudis, D., Jannopoulos, A., Flossos, A. & Zournatzidis, G. Monitoring the successful embolization of an arterio-venous fistula by a fibreoptic jugular vein catheter, 265
Petruzziello, L. *see* Agrò, F.
Philip, B.K., Pearman, M.H., Kovac, A. L., Chelly, J.E., Wetchler, B.V., McKenzie, R., Monk, T.G., Dershwitz, M., Mingus, M., Sung, Y.F., Hahne, W.F., Brown, R.A. & The Dolasetron PONV Prevention Study Group. Dolasetron for the prevention of postoperative nausea and vomiting following outpatient surgery with general anaesthesia: a randomized, placebo-controlled study, 23
Pippa, P. Brachial plexus block using a new subclavian perivascular technique: the proximal cranial needle approach, 120
Pitkänen, M. *see* Silvasti, M.
Podder, S., Wig, J., Malhotra, S.K. & Sharma, S. Effect of pre-emptive analgesia on self-reported and biological measures of pain after tonsillectomy, 319
Poelaert, J. *see* De Mey, J.C.
Pojer, H. *see* Krenn, H.
Popat, M.T., Chippa, J.H. & Russell, R. Awake fibreoptic intubation following failed regional anaesthesia for Caesarean section in a parturient with Still's disease, 211
Pourriat, J.-L. *see* Asehnoune, K.

Prien, T. *see* Buerkle, H.
Properzi, M. *see* Peduto, V.A.
Puehringer, F. *see* Kuenszberg, E.

Rae, C.P., Gallagher, G., Watson, S. & Kinsella, J. An audit of patient perception compared with medical and nursing staff estimation of pain during burn dressing changes, 43
Rangasami, J. *see* Shah, J.
Rathgeber, J. *see* Kazmaier, S.
Ravenscroft, P.M. *see* O'Riordan, J.
Redmond, H.P. *see* Goto, Y.
Reihnér, E., Grunditz, R., Giesecke, K. & Gustafsson, L.L. Postoperative nausea and vomiting after breast surgery: efficacy of prophylactic ondansetron and droperidol in a randomized placebo-controlled study, 197
Reinhart, K. *see* Meisner, M.
Reinhart, K., Sakka, S.G. & Meier-Hellmann, A. Haemodynamic management of a patient with septic shock, 6
Rodríguez, A. *see* Lucas, M.
Römer, A.W. *see* Azad, S.C.
Rosenberg, P.H. *see* Silvasti, M.
Rosseel, P.M.J. *see* El Azab, S.R.
Rubio, M. *see* Avellaneda, C.
Rumpold-Seitlinger, G. *see* Krenn, H.
Russell, R. *see* Popat, M.T.
Ruth, M. *see* Flucker, C.J.R.
Rutherford, A. *see* St Pierre, M.

Sağıroğlu, E. *see* Kızılarslan, S.
Saitoh, Y. *see* Fujii, Y.
Sakka, S.G. *see* Reinhart, K.
Sala, X. *see* Santos, P.
Sandström, K. *see* Bodelsson, G.
Sanjo, Y. *see* Suzuki, A.
Santos, P., Ayuso, A., Luis, M., Martínez, G. & Sala, X. Airway ignition during CO_2 laser laryngeal surgery and high frequency jet ventilation, 204
Sarmiento, J. *see* Avellaneda, C.
Sato, H. *see* Okuda, Y.
Sato, S. *see* Suzuki, A.
Satoh, T. *see* Adachi, Y.U.
Scandroglio, M. *see* Casati, A.
Scheffer, G.J. *see* El Azab, S.R.
Scherer, R. *see* Brodner, G.
Scherpereel, P. Foundation for European Education in Anaesthesiology, 75
Schlack, W. & Dinter, W. Haemodynamic effects of a left stellate ganglion block in ASA I patients. An echocardiographic study, 79
Scholz, J. *see* Martens, U.
Schöps, P. *see* Azad, S.C.
Schulte Am Esch, J. *see* Martens, U.
Seeberger, M.D. *see* Steiner, L.A.
Severijnen, R.S.V.M. *see* Brouwers, M.J.A.
Shah, J., Owen-Smith, O., Coley, S. & Rangasami, J. A survey of peripheral nerve stimulator (PNS) use, 273
Shah, M.V. *see* O'Connor, P.J.
Sharma, S. *see* Podder, S.
Shorten, G.D. *see* Goto, Y.
Shorten, G.D. *see* Halpenny, M.
Shorten, G.D. *see* Leonard, S.A.
Shorten, G.D. *see* Lydon, A.
Silvanus, M.-T. *see* Groeben, H.
Silvasti, M., Svartling, N., Pitkänen, M. & Rosenberg, P.H. Comparison of intravenous patient-controlled analgesia with tramadol versus morphine after microvascular breast reconstruction, 448
Skarvan, K. *see* Steiner, L.A.
Skelton, V.A., Henderson, K. & Liu, C. Anaesthetic implications of osteo-odonto-keratoprosthesis surgery, 390
Slappendel, R., Weber, E.W.G., Benraad, B., Van Limbeek, J. & Dirksen, R. Itching after intrathecal morphine. Incidence and treatment, 616
Smith, A., Pedersen, T. & Møller, A. The Cochrane Anaesthesia Review Group, 279
Snelgrove, H. & McAnulty, G. A glance at medical reading habits in Italy and England: is the writing on the wall?, 717
Snow, H.M. *see* Halpenny, M.
Sockalingham, I. *see* Gopinathan, C.
Sonntag, H. *see* Kazmaier, S.
St Pierre, M., Dunkel, M., Rutherford, A. & Hering, W. Does etomidate increase postoperative nausea? A double-blind controlled comparison

of etomidate in lipid emulsion with propofol for balanced anaesthesia, 634
Steinbeck, J. *see* Buerkle, H.
Steiner, L.A., Filipovic, M., Skarvan, K. & Seeberger, M.D. Anaesthetists as echocardiographers can influence the perioperative management of patients with suspected aortic stenosis, 463
Steinrücke, K. *see* Martens, U.
Stevens, A. *see* O'Riordan, J.
Stopp, N. *see* Jones, P.R.M.
Strobbe, J. *see* De Mey, J.C.
Sung, Y.F. *see* Philip, B.K.
Suzuki, A., Bito, H., Sanjo, Y., Katoh, T. & Sato, S. Evaluation of the PhysioFlex™ closed-circuit anaesthesia machine, 359
Svartling, N. *see* Silvasti, M.
Sweeney, B.P. *see* Hammond, E.J.
Szakmány, T. *see* Molnár, Z.

Tabib, A., Loire, R., Miras, A., Thivolet-Bejui, F., Timour, Q., Bui-Xuan, B. & Malicier, D. Unsuspected cardiac lesions associated with sudden unexpected perioperative death, 230
Taguchi, N. *see* Kihara, S.
Takahashi, S. *see* Yamaura, K.
Tanaka, H. *see* Fujii, Y.
Tanbo, T. *see* Bugge, J.F.
Tarantino, F. *see* Casati, A.
Tavernier, B. *see* Vallet, B.
Tebbs, S.E. *see* Moss, H.A.
Tekeres, M. *see* Molnár, Z.
Testa, C. *see* De Riu, P.L.
The Dolasetron PONV Prevention Study Group *see* Philip, B.K.
The Study Group Of Ketoprofen *see* Didailler, J.L.
Thivolet-Bejui, F. *see* Tabib, A.
Thoms, G. Anaesthesiology directors: more support on the way as Association of Anesthesia Clinical Directors goes international, 77
Timour, Q. *see* Tabib, A.
Tommasino, C. *see* Casati, A.
Törnebrandt, K. *see* Bodelsson, G.
Törnebrandt, K. *see* Flisberg, P.
Torri, G. *see* Casati, A.
Torri, G. *see* Casati, A.
Toyooka, H. *see* Fujii, Y.
Turgut, Ö. *see* Ceyhan, A.

Uchihashi, Y. *see* Adachi, Y.U.
Ünal, N. *see* Ceyhan, A.
Uyar, M. *see* Aysel, İ.

Valentini, G. *see* Agostoni, M.
Vallet, B., Tavernier, B. & Lund, N. Assessment of tissue oxygenation in the critically-ill, 221
Van Aken, H. *see* Baum, J.
Van Aken, H. *see* Brodner, G.
Van Aken, H. *see* Brodner, G.
Van Aken, H. *see* Buerkle, H.
Van Aken, H. Who should provide anaesthesia?, 535
Van Limbeek, J. *see* Slappendel, R.
Van Zalingen, E.E. *see* El-Kerdawy, H.M.M.
Veyckemans, F. *see* Pendeville, P.E.
Viby-Mogensen, J. Central anticholinergic syndrome or postoperative residual block?, 466
Vickers, M.D. Non-physician anaesthetists: can we agree on their role in Europe?, 537
Vislocký, I. *see* Cvachovec, K.
Von Montigny, S. *see* Pendeville, P.E.
Voyagis, G.S. & Dimitriou, V. Remifentanil vs. fentanyl during rigid bronchoscopy under general anaesthesia with controlled ventilation, 404
Voyagis, G.S. *see* Dimitriou, V.
Voyagis, G.S. *see* Dimitriou, V.

Wallerstedt, S.M. *see* Bodelsson, G.
Walsh, M. *see* Leonard, S.A.
Walther, B. *see* Flisberg, P.
Wang, C. *see* Pac-Soo, C.K.
Wang, J. *see* Goto, Y.
Wappler, F. *see* Martens, U.
Watanabe, K. *see* Adachi, Y.U.
Watanabe, S. *see* Kihara, S.
Watson, N.A., Beards, S.C., Altaf, N., Kassner, A. & Jackson, A. The effect of hyperoxia on cerebral blood flow: a study in healthy volunteers using magnetic resonance phase-contrast angiography, 152

Watson, S. *see* Rae, C.P.
Weber, E.W.G. *see* Slappendel, R.
Weigert, A. & Lawton, G. Accidental injection of thiopental into the epidural space, 69
Weissgarten, J. *see* Chanimov, M.
Weisz, M. *see* Flucker, C.J.R.
Welsh, K.R. *see* O'Connor, P.J.
Wetchler, B.V. *see* Philip, B.K.
Whitwam, J.G. *see* Pac-Soo, C.K.
Whitwam, J.G., Galletly, D.C., Ma, D. & Chakrabarti, M.K. The effects of propofol on heart rate, arterial pressure and Aδ and C somatosympathetic reflexes in anaesthetized dogs, 57
Wig, J. *see* Podder, S.
Wilkinson, D. *see* Kettler, D.
Wolfgart, M. *see* Buerkle, H.
Wouters, A. *see* De Wolff, M.H.

Yaguchi, Y. *see* Kihara, S.
Yamasaki, Y. *see* Kihara, S.
Yamaura, K., Higashi, M., Akiyoshi, K., Itonaga, Y., Inoue, H. & Takahashi, S. Pulmonary lipiodol embolism during transcatheter arterial chemoembolization for hepatoblastoma under general anaesthesia, 704

Zangrillo, A. *see* Agostoni, M.
Zeaki, M. *see* Michaloudis, D.
Zournatzidis, G. *see* Petru, A.

Keyword Index

ABDOMINAL AORTA
clamping, 491
ABNORMALITIES, MULTIPLE
Proteus syndrome, 645
ACID-BASE EQUILIBRIUM
metabolic acidosis, 256
ADRENERGIC ANTAGONISTS
clonidine, 379
ADRENERGIC ALPHA-ANTAGONISTS
clonidine, 379, 692
AIRWAY DEVICES
intubation tracheal, 46
technique, 46
training, 46
AIRWAY OBSTRUCTION, 398, 642
ALCOHOLS
ethanol, 559
ALLERGY AND IMMUNOLOGY, 248
clonidine, 50, 105
intravenous, 105
AMINO ACIDS
glycine, 559
ANAESTHESIA, 197, 443, 456, 591, 611
autonomic function, 99
autonomic nerve block, 79
brachial plexus, 566
caudal, 379
closed-circuit, 359
conduction, 348, 411, 566
diabetes mellitus, 99
early tracheal extubation, 111
epidural, 325, 379, 692
fibreoptic intubation, 211
gastroparesis, 99
general, 132, 230, 300, 306, 390, 411
intraocular pressure, 390
local, 379, 411
medical education, 91
medical informatics, 91
nerve block, 348, 566
obstetrics, 211, 542, 692
renal, 256
sciatic–femoral block, 348
spinal, 616
ANAESTHETIC TECHNIQUE, 185
balanced anaesthesia, 634
brachial plexus, 120
combined spinal epidural block, 329
continuous spinal anaesthesia, 239
epidural, 189, 542
general anaesthesia, 138
inhalational induction, 33
intravenous induction, 33
regional, 120, 542
ring block, 182
subcutaneous, 189
thoracic epidural, 627
ANAESTHETICS
desflurane, 325
etomidate, 248
fentanyl, 248
inhalation, 289, 359, 474, 506
intravenous, 57, 236, 248, 485
ketamine, 126, 248
lidocaine (lignocaine), 248
local, 348, 379, 436
midazolam, 248
propofol, 126, 138, 485
ropivacaine, 348
sevoflurane, 138
thiopental (thiopentone), 248, 485
ANAESTHETICS, INHALATION
isoflurane, 289, 506
nitrous oxide, 359
sevoflurane, 359, 474, 662
ANAESTHETICS, INTRAVENOUS
complications, 236
etomidate, 248
fentanyl, 248
ketamine, 126, 248
midazolam, 248
propofol, 57, 126, 485, 720
thiopental (thiopentone), 248, 485
ANAESTHETICS, LOCAL
alkalinization, 38
benzalkonium compounds, 680
bupivacaine, 189, 239, 379, 627, 692
levobupivacine, 373
lidocaine (lignocaine), 248, 436, 672
mepivacaine, 627

© 2000 European Academy of Anaesthesiology

ropivacaine, 325, 348
sodium bicarbonate, 38

ANALGESIA
bupivacaine, 319
epidural, 132, 566, 751
intercostal nerve block, 132
intra-articular, 295
intravenous, 566
intravenous opioids, 448
patient-controlled, 431, 566
peripheral, 295
postoperative, 239, 448
pre-emptive analgesia, 319

ANALGESICS
acetaminophen (paracetamol), 576
clonidine, 379
diamorphine, 189
fentanyl, 348, 692
ketorolac, 85
metamizol, 85
morphine, 448, 611
opioid, 189, 348, 379, 448, 751
propacetamol, 85
sufentanil, 379
tramadol, 448, 576

ANEURYSM
aortic dissection, 709

ANOXIA
anoxaemia, 456

ANTIEMETICS
5-HT_3 antagonist, 23
droperidol, 611
dolasetron, 23
emesis, 23
nausea, 23
ondansetron, 611

ANTI-INFECTIVE AGENTS, 680

ARTERIES
aorta, 709
coronary vessels, 481, 485
omental, 720
thoracic, 709

ARTHROPLASTY
hip, 431
knee, 431

ATMOSPHERE
air, 512

AUDIT
analgesia, 43
pain, 43

AUTONOMIC NERVOUS SYSTEM
hyper-reflexia, 329
paraplegia, 329

BACTERIAL INFECTIONS
gram-positive bacterial infections, 289
gram-negative bacterial infections, 289

BENZODIAZEPINES
midazolam, 126

BISPECTRAL INDEX, 499

BLOOD
chemical analysis, 132

BLOOD COAGULATION (*see also* COAGULATION)
fibrinolysis, 173

BLOOD COAGULATION TESTS (*see also* COAGULATION)
cardiopulmonary bypass, 173
thromboelastography, 173

BLOOD TRANSFUSION
autologous, 411

BODY POSITION
Trendelenburg, 300

BREAST SURGERY, 197

BRONCHIAL DISEASES
bronchial fistula, 418
bronchial spasm, 672

BURNS
analgesia, 43

BUTYROPHENONES
droperidol, 126

CAESAREAN SECTION, 329

CALCIUM CHANNEL BLOCKERS
clevidipine, 506
nimodipine, 383

CARBON DIOXIDE, 622
arterial, 306
gradients, 306
insufflation, 256
tension, 306

CARDIOPULMONARY BYPASS, 499

CARDIOVASCULAR PHYSIOLOGY
haemodynamics, 79
ventricular function, left, 79

CARDIOVASCULAR SYSTEM
effects, 168
CATECHOLAMINES, 168
CATHETERIZATION, 680
adverse effects, 168
central venous, 168
CELL DEATH
apoptosis, 474
CENTRAL NERVOUS SYSTEM AGENTS
clonidine, 692
CENTRAL VENOUS, 680
CEREBROVASCULAR CIRCULATION
autoregulation, 152
cerebral blood flow, 152
CIRCULATION
heart rate and blood pressure, 57
CLONIDINE
epidural, 50
intravenous, 50
COAGULATION (*see also* BLOOD COAGULATION)
clotting effects, 373
thromboelastography, 373
COMPLEX REGIONAL PAIN SYNDROME
axillary brachial plexus catheter, 185
morphine, 185
COMPLIANCE
atropine, 236
bronchospasm, 236
COMPLICATIONS, 431
nausea, 64
vomiting, 64
CONTRAST MEDIA
iodized oil, 704
CORONARY VESSELS, 485

DATA COLLECTION
computer, 284
interviews, 284
DEATH
cardiac, 230
sudden, 230
DISSECTING
aortic dissection, 709
DISSOCIATIVE
ketamine, 126
DIURETICS
mannitol, 559
DRUG DELIVERY, 160
DRUG INTERACTIONS, 126
DRUG SYNERGISM
droperidol, 126
midazolam, 126
propofol, 126
DRUGS
antiemetics, 197
clonidine, 295
droperidol, 197
morphine, 295
ondansetron, 197

ECHOCARDIOGRAPHY
transoesophageal, 709
EDUCATION, 443
ELECTROCARDIOGRAPHY
epinephrine, 132
potassium, 132
QT-interval, 662
ELECTROENCEPHALOGRAPHY, 105, 499
EMBOLISM
pulmonary embolism, 704
ENDOTHELIUM, 481, 485
vascular, 720
ENTEROBACTERIACEAE INFECTIONS
Escherichia coli infections, 289
EPIDURAL
anaesthesia, 379
caudal, 379
morphine, 627
EPIDURAL ANAESTHESIA
clonidine, 50
morphine, 50
EQUIPMENT AND SUPPLIES, 456
mechanical, 354
ventilators, 354
EQUIPMENT CONTAMINATION, 512
EQUIPMENT FAILURE, 456, 591
EQUIPMENT HAZARD, 512
EQUIPMENT SAFETY, 456
ETILEFRINE
hypotension, 177
spinal anaesthesia, 177
EXTRACORPOREAL CIRCULATION
coronary artery bypass, 601

EYE
anterior eye segment, 390
cornea, 390

FAILURE, RENAL, 491
FENOLDOPAM, 491
FISTULA
direct surgical embolization, 265
post-traumatic carotid-cavernous fistula, 265
surgical repair, 265

GASES
oxygen, 456, 591
GASTRO-OESOPHAGEAL REFLUX, 354
GENERAL ANAESTHESIA
obesity, 300
pneumoperitoneum, 300
GRAM-POSITIVE BACTERIAL INFECTIONS
staphylococcal infections, 289
GRAM-NEGATIVE BACTERIAL INFECTIONS
Enterobacteriaceae infections, 289
GYNAECOLOGICAL, 611

HAEMODIALYSIS
haemodiafiltration, 665
HAEMODYNAMICS
blood pressure, 744
HAEMOSTASIS, 173
HAMARTOMA
Proteus syndrome, 645
HEALTH CARE COSTS
hospital costs, 566
HEART
bundle of His, 230
heart conduction system, 230
myocardium, 230
HEART SURGERY, 85
HETEROCYCLIC COMPOUNDS
etomidate, 634
HIGH DEPENDENCY, 111
HIGH PRESSURE LIQUID CHROMATOGRAPHY, 18
HIRSCHSPRUNG DISEASE, 751
HOMEOSTASIS
water-electrolyte balance, 559

IMIDAZOLES
etomidate, 634
IMMUNE SYSTEM
neutrophils, 474
phagocytes, 474
IMMUNITY
neutrophil activation, 474
INFECTION
sepsis, 665, 680
septicaemia, 680
INFLAMMATION, 173
INFLAMMATORY MEDIATORS
chemokines, 474
INHALATIONAL ANAESTHETICS
desflurane, 311
enflurane, 311
halothane, 311
isoflurane, 289, 311, 506
nitrous oxide, 311
sevoflurane, 662
INTENSIVE CARE, 111
INTERVENTIONAL, 704
INTRAOPERATIVE
pulse oximetry, 704
INTRAOPERATIVE COMPLICATIONS, 230
pulmonary embolism, 704
INTRATHECAL OPIOIDS
fentanyl, 239
INTRATRACHEAL, 236, 587
laryngeal masks, 354, 642, 744
INTUBATION, 587
intratracheal, 395, 436, 744
laryngeal masks, 354, 395, 642

LARYNGEAL CARTILAGES
cricoid, 443
LARYNX, 443
LASER SURGERY
airway fire, 204
high frequency jet ventilation, 204
LEUKOCYTES MONONUCLEAR
lymphocytes, 248
monocytes, 248

MALIGNANT HYPERTHERMIA
caffeine, 364

halothane, 364
inositol 1, 4, 5-trisphosphate, 364
in vitro contracture test, 364
mononuclear white blood cells, 364
ryanodine, 364
MEASUREMENT
body height, 698
body weight, 698
MEDICAL
paediatrics, 379
METABOLISM, 221
MINIMAL ALVEOLAR CONCENTRATION
isoflurane, 105
MONITORING, 622
intraoperative, 359, 583
physiological, 168
pulse oximetry, 704
MUSCLE, SKELETAL
adductor pollicis, 583
masseter, 583
MYOCARDIAL DISEASE
cardiomyopathy, 230
MYOCARDIAL ISCHAEMIA
coronary disease, 601

NARCOTIC ANTAGONISTS
naloxone, 616
NERVE BLOCK
epidural, 373
NERVE FIBRE
Aδ and C, 57
NERVES
renal sympathetic nerves, 311
NEUROMUSCULAR BLOCKADE, 583
NEUROMUSCULAR BLOCKING AGENTS, 583
NEUROMUSCULAR JUNCTION DISEASES
myasthenia gravis, 325
NEUROMUSCULAR NON-DEPOLARIZING AGENTS
atracurium, 481
cisatracurium, 583
pancuronium, 481
rocuronium, 481
vecuronium, 383, 481
NITRIC ACID, 160
NITRIC OXIDE
pregnancy, 269
NITRIC OXIDE ADMINISTRATION
lung model, 160
NITROGEN DIOXIDE, 160
NITROUS OXIDE
legal limits, 149
staff exposure, 149
NOCICEPTIVE REFLEX, 57
NSAIDS
tenoxicam, 239
NUCLEAR MAGNETIC RESONANCE
cerebral blood flow, 152

OBESITY
cardiopulmonary function, 300
OESOPHAGEAL ACHALASIA, 398
OPHTHALMOLOGICAL SURGICAL PROCEDURES
cataract extraction, 474
OPIOIDS
fentanyl, 692
morphine, 448, 616, 627, 751
remifentanil, 111
sufentanil, 379
tramadol, 448, 576
ORGANIC CHEMICALS
propofol, 634
ORTHOPAEDIC PROCEDURES
arthoplasty, 411
arthoplasty replacement, 411
hip, 411
knee, 411
replacement, 411
ORTHOPAEDICS, 411
OSMOTIC
mannitol, 559
OTORHINOLARYNGOLOGICAL SURGICAL PROCEDURES
tonsillectomy, 576
OUTPATIENT SURGERY, 23
OXYGEN DEMAND, 221
OXYGEN MONITORING
fibreoptic jugular vein oxygen saturation catheter, 265
intravascular catheter, 265
jugular vein oxygen monitoring, 265
OXYGEN SUPPLY, 221

PAEDIATRICS, 576

PAIN, 448
EMLA, 182
postoperative, 189, 627
PAIN MANAGEMENT, 185
PAIN MEASUREMENT
child, 284
PATHOLOGICAL PROCESSES
sepsis, 665
PATIENT-CONTROLLED ANALGESIA
postoperative, 448
PATIENTS
high-risk elderly, 239
PHARMACOLOGY
drug interactions, 383
propofol, 499
sufentanil, 499
PHARYNGITIS, 436
PHENOLS
propofol, 634
PHYSIOLOGICAL, 622
POSTOPERATIVE COMPLICATIONS, 173
postoperative nausea and vomiting, 23, 197, 261, 448, 611, 634
POSTOPERATIVE PAIN, 85
PROPOFOL
biodisposition, 18
red blood cells, 18
PROSTATECTOMY
transurethral resection of prostate, 559
PROTHESES AND IMPLANTS
intraocular, 390
lenses, 390
osteo-odontokeratoprosthesis, 390

RABBIT, 18
RADIOLOGY, 704
RECEPTOR AGONIST
dopamine, 491
RECOVERY PERIOD
ambulatory surgery, 138
outpatients, 138
REGIONAL ANAESTHESIA
blood patch, 208
epidural blockade, 208
stellate ganglion, 79
REGIONAL BLOOD FLOW
oxygen saturation, 152
RENAL TRANSPLANTATION
diabetes, 729
haemodynamics, 729
uraemia, 729
RESPIRATION, ARTIFICIAL, 601, 622
high-frequency jet ventilation, 418
high-frequency ventilation, 418
intermittent positive-pressure ventilation, 306
positive-pressure ventilation, 737
RESPIRATORY HYPERSENSITIVITY
asthma, 672
RESPIRATORY INSUFFICIENCY, 418
RESPIRATORY MECHANICS
aspiration, 354
respiration, 354
RESPIRATORY THERAPY, 601

SEPSIS, 6
SHOCK
haemodynamic management, 6
SKIN DISEASES
pruritus, 616
SLEEP APNOEA SYNDROMES
sleep apnoea, 751
central, 751
SPECIALTIES, 411
medical, 379
surgical, 411
SPINAL, 616
SURGERY
abdominal, 111
breast, 261
ingrowing toenail, 182
laparoscopic cholecystectomy, 64
liver transplantation, 111
microvascular breast reconstruction, 448
retroperitoneoscopy, 256
upper abdominal, 627
SURGICAL,
orthopaedics, 411
SYMPATHETIC NERVOUS SYSTEM, 57

TACHYKININS
substance P, 720
TEACHING
patient simulation, 443

THORACIC SURGERY
catecholamines, 50
cortisol β-endorphin, 50
leucocytes, 50
stress responses, 50
vasopressin, 50
TISSUE CULTURE
cell culture, 248
TONSILLECTOMY, 319
TRACHEA
tracheomalacia, 642
TRACHEOSTOMY
percutaneous, 688

ULTRASONOGRAPHY
echocardiography, 79
URINARY RETENTION, 431
URINARY TRACT
bladder, 559

VARICELLA PNEUMONIA, 269
VEINS
omental, 720
VENTILATORS
mechanical, 359, 737
VOMITING
antiemetics, 64
combination, 64
dexamethasone, 64
granisetron, 64

WATER, 512

Subject Index

A central venous catheter coated with benzalkonium chloride for the prevention of catheter-related microbial colonization (Moss *et al.*) 2000 **17**, 680–687

A comparative study of patient-controlled epidural diamorphine, subcutaneous diamorphine and an epidural diamorphine/bupivacaine combination for postoperative pain (Gopinathan *et al.*) 2000 **17**, 189–196

A comparison of the effects on postoperative pain relief of epidural analgesia started before or after surgery (Flisberg *et al.*) 2000 **17**, 627–633

A retrospective survey of fibrinolysis as an indicator of poor outcome after cardiopulmonary bypass and a possible early sign of systemic inflammation syndrome (Cvachovec *et al.*) 2000 **17**, 173–176

Abstracts

, EAA Meeting Abstracts, September 2000, Paris 2000 **17**, 754–785

, Symposium: Simulation in Anaesthesia and Intensive Care Medicine 2000. Annual meeting of the 'Society in Europe for Simulation Applied to Medicine' (SESAM) 2000 **17**, 515–526

Accuracy of end-tidal carbon dioxide monitoring using the NBP-75® microstream capnometer. A study in intubated ventilated and spontaneously breathing nonintubated patients (Casati *et al.*) 2000 **17**, 622–626

Acid-base equilibrium during capnoretroperitoneoscopic nephrectomy in patients with end-stage renal failure: a preliminary report (Demian *et al.*) 2000 **17**, 256–260

Acute pain management: analysis, implications and consequences after prospective experience with 6349 surgical patients (Brodner *et al.*) 2000 **17**, 566–575

Alkalinization of amide local anaesthetics by addition of 1% sodium bicarbonate solution (Milner *et al.*) 2000 **17**, 38–42

An audit of patient perception compared with medical and nursing staff estimation of pain during burn dressing changes (Rae *et al.*) 2000 **17**, 43–45

Anaesthetic implications of osteo-odonto-keratoprosthesis surgery (Skelton *et al.*) 2000 **17**, 390–394

Are diabetic patients in danger at renal transplantation? An invasive perioperative study (Norio *et al.*) 2000 **17**, 729–736

Ballotability of cuff to confirm the correct intratracheal position of the endotracheal tube in the intensive care unit (Pattnaik & Bodra) 2000 **17**, 587–590

Bedside percutaneous tracheostomy experience with 72 critically ill patients (Kahveci *et al.*) 2000 **17**, 688–691

Book reviews

, Anaesthesia for Obstetrics and Gynaecology (Russell) 2000 **17**, 714

, Anaesthesia for Vascular Surgery (Bannister & Wildsmith) 2000 **17**, 469–470

, Cardiovascular Drugs in the Perioperative Period (Foëx, Harrison & Opie) 2000 **17**, 654

, Clinical Pharmacology of Local Anaesthetics (Tetzlaff) 2000 **17**, 278

, Critical Care Focus Volume 2: Respiratory Failure (Galley) 2000 **17**, 788

, Final FRCA Multiple Choice Questions (Brunner, Robinson & Williams) 2000 **17**, 531

, Fundamentals of Anaesthesia (Pinnock, Lin, Smith & Jones) 2000 **17**, 468

, Fundamentals of Operating Department Practice (Davey & Ince) 2000 **17**, 714

, Lee's Synopsis of Anaesthesia (Rushman, Davies & Cashman) 2000 **17**, 215

, Nunn's Applied Respiratory Physiology (Lumb) 2000 **17**, 339–340

, Pharmacology for Anaesthesia and Intensive Care (Peck & Williams) 2000 **17**, 531

, Principles and Practice of Obstetric Anaesthesia and Analgesia (Holdcroft & Thomas) 2000 **17**, 406

, The Effective Management of Post-Operative Nausea and Vomiting (Strunin, Rowbotham & Miles) 2000 **17**, 215–216

, Thoracic Anaesthesia: Principles and Practice (Ghosh & Latim) 2000 **17**, 339

Brachial plexus block using a new subclavian perivascular technique: the proximal cranial needle approach (Pippa) 2000 **17**, 120–125

Can computer-based assessment help us understand children's pain? (Calam *et al.*) 2000 **17**, 284–288

Effects of isoflurane on bacterial growth (Asehnoune *et al.*) 2000 **17**, 289–294

Can we improve the estimate of arterial PCO_2 from end-tidal PCO_2? (Fletcher & Boris-Möller) 2000 **17**, 306–310

Cardiovascular responses to pulmonary artery catheterization (O'Connor *et al.*) 2000 **17**, 168–172

© 2000 European Academy of Anaesthesiology

Clinical letters

, Airway ignition during CO_2 laser laryngeal surgery and high frequency jet ventilation (Santos *et al.*) 2000 **17**, 204–207

, Airway obstruction during general anaesthesia in a child with congenital tracheomalacia (Okuda *et al.*) 2000 **17**, 642–644

, Anaesthesia for Proteus syndrome (Ceyhan *et al.*) 2000 **17**, 645–647

, Awake fibreoptic intubation following failed regional anaesthesia for Caesarean section in a parturient with Still's disease (Popat *et al.*) 2000 **17**, 211–214

, Combination of balanced and regional anaesthesia for minimally invasive surgery in a patient with myasthenia gravis (Hübler *et al.*) 2000 **17**, 325–328

, Combined spinal–epidural anaesthesia for Caesarean section in a paraplegic woman: difficulty in obtaining the expected level of block (Agostoni *et al.*) 2000 **17**, 329–331

, Contamination of piped medical gas supply with water (Hay) 2000 **17**, 512–514

, Delivery of an hypoxic gas mixture due to a defective rubber seal of a flowmeter control tube (Hay) 2000 **17**, 456–458

, Epidural analgesia in a newborn with Hirschsprung's disease, associated with congenital central hypoventilation syndrome (Brouwers *et al.*) 2000 **17**, 751–753

, Monitoring the successful embolization of an arterio-venous fistula by a fibreoptic jugular vein catheter (Petru *et al.*) 2000 **17**, 265–268

, Near-fatal acute dissection of the thoracic aorta involving the right coronary artery (Chevalley *et al.*) 2000 **17**, 709–711

, Nitric oxide in the treatment of fulminant pulmonary failure in a young pregnant woman with varicella pneumonia (Bugge & Tanbo) 2000 **17**, 269–272

, Oxygen delivery failure resulting from interference with a Bain breathing system (Hay) 2000 **17**, 591–593

, Pulmonary lipiodol embolism during transcatheter arterial chemoembolization for hepatoblastoma under general anaesthesia (Yamaura *et al.*) 2000 **17**, 704–708

, Rapidly developing airway obstruction resulting from achalasia of the oesophagus (Ray) 2000 **17**, 398–400

, Successful epidural blood patch in a patient with headache for 6 months after lumbar root decompression (Gentili *et al.*) 2000 **17**, 208–210

, Use of the intubating laryngeal mask for airway management and light-guided tracheal intubation in the lateral position (Dimitriou & Voyagis) 2000 **17**, 395–397

Combined sciatic–femoral nerve block with 0.75% ropivacaine: effects of adding a systemically inactive dose of fentanyl (Magistris *et al.*) 2000 **17**, 348–353

Comparison of cisatracurium-induced neuromuscular block at the masseter and adductor pollicis muscle (de Rossi *et al.*) 2000 **17**, 583–586

Comparison of intravenous patient-controlled analgesia with tramadol versus morphine after microvascular breast reconstruction (Silvasti *et al.*) 2000 **17**, 448–455

Comparison of the effects of inhalational anaesthetic agents on sympathetic activity in rabbits (Pac-Soo *et al.*) 2000 **17**, 311–318

Comparison of ventilatory and haemodynamic effects of BIPAP and S-IMV/PSV for postoperative short-term ventilation in patients after coronary artery bypass grafting (Kazmaier *et al.*) 2000 **17**, 601–610

Continuous axillary brachial plexus analgesia with low dose morphine in patients with complex regional pain syndromes (Azad *et al.*) 2000 **17**, 185–188

Continuous spinal anaesthesia/analgesia for the perioperative management of high-risk patients (Michaloudis *et al.*) 2000 **17**, 239–247

Correspondence

, 'Diprivan': a change of formulation (Hart) 2000 **17**, 71–73

, A rare misplacement of a central venous catheter (Khan & Khan) 2000 **17**, 527–528

, A successfully resuscitated case of amniotic fluid embolism (Aziz) 2000 **17**, 712–713

, A survey of peripheral nerve stimulator (PNS) use (Shah *et al.*) 2000 **17**, 273–275

, Accidental injection of thiopental into the epidural space (Weigert & Lawton) 2000 **17**, 69–70

, Anaesthesia in lipid myopathy (Lucas *et al.*) 2000 **17**, 461–462

, Anaesthetists as echocardiographers can influence the perioperative management of patients with suspected aortic stenosis (Steiner *et al.*) 2000 **17**, 463

, Bradycardia, hypotension and cerebral dysfunction after spinal anaesthesia (Payne) 2000 **17**, 334–335

, Central anticholinergic syndrome or postoperative residual block? (Viby-Mogensen) 2000 **17**, 466

, Continuous spinal anaesthesia or continuous epidural anaesthesia for postoperative pain con-

trol after hip replacement (Brodner *et al.*) 2000 **17**, 402–403
, Effects of a pneuperitoneum in the obese patient (Dumont & Mardirosoff) 2000 **17**, 786
, Gastroscopy in awake and anaesthetized patients using a modified laryngeal mask (Agrò *et al.*) 2000 **17**, 652–653
, High subarachnoid block for severe bronchospasm (Pakulski *et al.*) 2000 **17**, 594–595
, Induction and maintenance of anaesthesia with sevoflurane in comparison to high dose opioid during coronary artery bypass surgery (Azab *et al.*) 2000 **17**, 336–338
, Knowledge about pulse oximetry among residents and nurses (Bilgin *et al.*) 2000 **17**, 650–651
, Microalbuminuria and serum procalcitonin levels following oesophagectomy (Molnár *et al.*) 2000 **17**, 464–465
, Prognostic value of Apache II and III for preoperative evaluation of emergency surgery patients (Krenn *et al.*) 2000 **17**, 529–530
, Recurarization in the recovery room (Milliet) 2000 **17**, 332
, Remifentanil vs. fentanyl during rigid bronchoscopy under general anaesthesia with controlled ventilation (Voyagis & Dimitriou) 2000 **17**, 404–405
, The anaesthetic potency of propofol in the rat (Crozier) 2000 **17**, 276
, The analgesic efficacy and tolerance of ketoprofen (100 mg) combined with morphine in patient-controlled analgesia after orthopaedic surgery (Didailler *et al.*) 2000 **17**, 459–460
, The psychology of human error revisited (Klemola) 2000 **17**, 401
, Who should provide anaesthesia? (Lassner) 2000 **17**, 648–649

Disposition of propofol between red blood cells, plasma, brain and cerebrospinal fluid in rabbits (De Riu *et al.*) 2000 **17**, 18–22
Do muscle relaxants influence vascular tone in isolated coronary artery segments? (Klockgether-Radke *et al.*) 2000 **17**, 481–484
Does etomidate increase postoperative nausea? A double-blind controlled comparison of etomidate in lipid emulsion with propofol for balanced anaesthesia (St Pierre *et al.*) 2000 **17**, 634–641
Dolasetron for the prevention of postoperative nausea and vomiting following outpatient surgery with general anaesthesia: a randomized, placebo-controlled study (Philip *et al.*) 2000 **17**, 23–32
Double-blind randomized study of tramadol vs. paracetamol in analgesia after day-case tonsillectomy in children (Pendeville *et al.*) 2000 **17**, 576–582

Editorials
, Anaesthesia organizations in Europe: a coming together (Adams) 2000 **17**, 1
, Anaesthesiology directors: more support on the way as Association of Anesthesia Clinical Directors goes international (Thoms) 2000 **17**, 77–78
, Calcium hydroxide lime – a new carbon dioxide absorbent: a rationale for judicious use of different absorbents (Baum & Van Aken) 2000 **17**, 597–600
, Clinical governance (Fletcher) 2000 **17**, 471–473
, Confederation of European National Societies of Anaesthesiology (CENSA) (Kettler & Wilkinson) 2000 **17**, 145
, Foundation for European Education in Anaesthesiology (Scherpereel) 2000 **17**, 75–76
, Opioid detoxification under anaesthesia (Bovill) 2000 **17**, 657–661
, Preoperative fasting and administration of regular medications in adult patients presenting for elective surgery. Has new evidence changed clinical practice? (Connolly & Cunningham) 2000 **17**, 219–220
, Procedure coding in anaesthesia (Banks) 2000 **17**, 407–410
, Questions about the Macintosh laryngoscope and technique of laryngoscopy (Henderson) 2000 **17**, 2–5
, The Cochrane Anaesthesia Review Group (Smith *et al.*) 2000 **17**, 279–283
, The European Society of Anaesthesiologists and the EJA: strength in unity (Aitkenhead) 2000 **17**, 217–218
, Thromboembolism and thromboprophylaxis in high risk surgery: facts and assumptions – a topic for emotions? (Dahl) 2000 **17**, 343–347
, What does the bispectral EEG index monitor? (Chan & Gin) 2000 **17**, 146–148
, Who should provide anaesthesia? (Van Aken) 2000 **17**, 535–536

Effect of clevidipine, an ultra-short acting 1,4-dihydropyridine calcium channel blocking drug, on the potency of isoflurane in rats and dogs (de Wolff *et al.*) 2000 **17**, 506–511
Effect of epidural and intravenous clonidine on the neuro-endocrine and immune stress response in

patients undergoing lung surgery (Novak-Jankovič *et al.*) 2000 **17**, 50–56
Effect of pre-emptive analgesia on self-reported and biological measures of pain after tonsillectomy (Podder *et al.*) 2000 **17**, 319–324
Effects of normothermic cardiopulmonary bypass on bispectral index (Hirschi *et al.*) 2000 **17**, 499–505
Effects of pneumoperitoneum and reverse Trendelenburg position on cardiopulmonary function in morbidly obese patients receiving laparoscopic gastric banding (Casati *et al.*) 2000 **17**, 300–305
Effects of propofol on substance P-induced relaxation in isolated human omental arteries and veins (Bodelsson *et al.*) 2000 **17**, 720–728
Efficacy of prophylactic droperidol, ondansetron or both in the prevention of postoperative nausea and vomiting in major gynaecological surgery. A prospective, randomized, double-blind clinical trial (Peixoto *et al.*) 2000 **17**, 611–615
Electronic data collection by trainee anaesthetists using palm top computers (Hammond & Sweeney) 2000 **17**, 91–98
Elimination of procalcitonin and plasma concentrations during continuous veno-venous haemodiafiltration in septic patients (Meisner *et al.*) 2000 **17**, 665–671
Epidural fentanyl–bupivacaine compared with clonidine–bupivacaine for analgesia in labour Kızılarslan *et al.*) 2000 **17**, 692–697
Epinephrine, potassium and the electrocardiogram during regional anaesthesia (Hahn & Löfgren) 2000 **17**, 132–137
Erratum
, European Society of Anaesthesiologists' Abstracts, Supplement 19. Abstract A-604, Doepfmer UR *et al.* 2000 **17**, 655

Evaluation of standard haemodynamic tests of autonomic function and HbA1c as predictors of delayed gastric emptying in patients with type 1 diabetes mellitus (Lydon *et al.*) 2000 **17**, 99–104
Evaluation of the effects of levobupivacaine on clotting and fibrinolysis using thromboelastography (Leonard *et al.*) 2000 **17**, 373–378
Evaluation of the PhysioFlex™ closed-circuit anaesthesia machine (Suzuki *et al.*) 2000 **17**, 359–363
Evaluation of the VersaMed portable ventilator: clinical trials (Brill *et al.*) 2000 **17**, 737–743

General versus regional anaesthesia for cataract surgery: effects on neutrophil apoptosis and the postoperative pro-inflammatory state (Goto *et al.*) 2000 **17**, 474–480
Granisetron/dexamethasone combination for the prevention of postoperative nausea and vomiting after laparoscopic cholecystectomy (Fujii *et al.*) 2000 **17**, 64–68

Haemodynamic effects of a left stellate ganglion block in ASA I patients. An echocardiographic study (Schlack & Dinter) 2000 **17**, 79–84
Haemodynamic responses to the intubating laryngeal mask and timing of removal (Kihara *et al.*) 2000 **17**, 744–750
High-frequency jet ventilation in European and North American institutions: developments and clinical practice (Ihra *et al.*) 2000 **17**, 418–430

Incidence and risk factors for allogenic blood transfusion during major joint replacement using an integrated autotransfusion regimen (Borghi & Casati) 2000 **17**, 411–417
Inhalational versus intravenous induction. A survey of emergency anaesthetic practice in the United Kingdom (Moore & Davies) 2000 **17**, 33–37
Inositol 1,4,5-trisphosphate synthesis in mononuclear white blood cells of malignant hypether following in vitro exposure to halothane, caffeine and ryanodine (Martens *et al.*) 2000 **17**, 364–372
Intra-articular clonidine analgesia after knee arthroscopy (Buerkle *et al.*) 2000 **17**, 295–299
Itching after intrathecal morphine. Incidence and treatment (Slappendel *et al.*) 2000 **17**, 616–621

Lidocaine 10% in the endotracheal tube cuff: blood concentrations, haemodynamic and clinical effects (Altinaş *et al.*) 2000 **17**, 436–442
Lidocaine inhalation for local anaesthesia and attenuation of bronchial hyper-reactivity with least airway irritation. Effect of three different dose regimens (Groeben *et al.*) 2000 **17**, 672–679

Modification of the Penlon Nuffield 200 series ventilator and IPPV with a laryngeal mask airway (Miller) 2000 **17**, 354–358

Nitric oxide delivery to the lung: a model (Jones *et al.*) 2000 **17**, 160–167

Patient-controlled analgesia and urinary retention following lower limb joint replacement: prospec-

tive audit and logistic regression analysis (O'Riordan *et al.*) 2000 **17**, 431–435

Personal view

, Non-physicians anaesthetists: can we agree on their role in Europe? (Vickers) 2000 **17**, 537–541

Postoperative nausea and vomiting after breast surgery: efficacy of prophylactic ondansetron and droperidol in a randomized placebo-controlled study (Reihnér *et al.*) 2000 **17**, 197–203

Postoperative nausea and vomiting in women following breast surgery: an audit (Jaffe *et al.*) 2000 **17**, 261–264

Propofol and thiopental attenuate the contractile response to vasoconstrictors in human and porcine coronary artery segments (Klockgether-Radke *et al.*) 2000 **17**, 485–490

Reducing the demand for admission to intensive care after major abdominal surgery by a change in anaesthetic practice and the use of remifentanil (Park *et al.*) 2000 **17**, 111–119

Reviews

, Assessment of tissue oxygenation in the critically-ill (Vallet *et al.*) 2000 **17**, 221–229

, Epidural analgesia–anaesthesia in obstetrics (Gomar & Fernandez) 2000 **17**, 542–558

, Haemodynamic management of a patient with septic shock (Reinhart *et al.*) 2000 **17**, 6–17

Sevoflurane progressively prolongs the QT interval in unpremedicated female adults (Kuenszberg *et al.*) 2000 **17**, 662–664

Sevoflurane provides better recovery than propofol plus fentanyl in anaesthesia for day-care surgery (Peduto *et al.*) 2000 **17**, 138–143

Small dose midazolam or droperidol reduces the hypnotic dose of propofol at the induction of anaesthesia (Adachi *et al.*) 2000 **17**, 126–131

Staff exposure to anaesthetic gases in theatre and non theatre areas (Henderson & Matthews) 2000 **17**, 149–151

Substances used for local and general anaesthesia in major surgery suppress proliferative responsiveness of normal rat peripheral blood mononuclear cells in culture (Chanimov *et al.*) 2000 **17**, 248–255

The 50-millilitre syringe as an inexpensive training aid in the application of cricoid pressure (Flucker *et al.*) 2000 **17**, 443–447

The accuracy of the estimation of body weight and height in the intensive care unit (Leary *et al.*) 2000 **17**, 698–703

The effect of a single intravenous dose of metamizol 2 g, ketorolac 30 mg, and propacetamol 1 g on haemodynamic parameters and postoperative pain after heart surgery (Avellaneda *et al.*) 2000 **17**, 85–90

The effect of hyperoxia on cerebral blood flow: a study in healthy volunteers using magnetic resonance phase-contrast angiography (Watson *et al.*) 2000 **17**, 152–159

The effect of oral etilefrine premedication on the incidence of hypotension during spinal anaesthesia (Bouaggad *et al.*) 2000 **17**, 177–181

The effects of atropine on dynamic compliance in healthy intubated adults (Nielsen *et al.*) 2000 **17**, 236–238

The effects of fenoldopam on renal blood flow and tubular function during aortic cross-clamping in anaesthetized dogs (Halpenny *et al.*) 2000 **17**, 491–498

The effects of nimodipine on vecuronium-induced neuromuscular blockade (Aysel *et al.*) 2000 **17**, 383–389

The effects of propofol on heart rate, arterial pressure and Aδ and C somatosympathetic reflexes in anaesthetized dogs (Whitwam *et al.*) 2000 **17**, 57–63

The influence of sufentanil and/or clonidine on the duration of analgesia after a caudal block for hypospadias repair surgery in children (De Mey *et al.*) 2000 **17**, 379–382

The influence of the α_2- adrenoceptor agonist, clonidine, on the EEG and on the MAC of isoflurane (El-Kerdawy *et al.*) 2000 **17**, 105–110

The use of EMLA reduces the pain associated with digital ring block for ingrowing toenail correction (Browne *et al.*) 2000 **17**, 182–184

The volumetric fluid balance as a measure of fluid absorption during transurethral resection of the prostate (Hahn) 2000 **17**, 559–565

Unsuspected cardiac lesions associated with sudden unexpected perioperative death (Tabib *et al.*) 2000 **17**, 230–235

Use of an illuminated flexible catheter for light-guided tracheal intubation through the intubating laryngeal mask by nurses (Dimitriou & Voyagis) 2000 **17**, 46–49